

Enstilar[®]
(calcipotriene and betamethasone
dipropionate) Foam 0.005%/0.064%

Treating plaque psoriasis with Enstilar[®] Spray Foam

**SEE OUR SAVINGS
OFFER INSIDE**

Not an
actual patient.

IMPORTANT SAFETY INFORMATION

What is Enstilar Foam?

Enstilar[®] (calcipotriene and betamethasone dipropionate) Foam is a prescription medicine used on the skin (topical) to treat plaque psoriasis in people 12 years of age and older. It is not known if Enstilar Foam is safe and effective in children under 12 years of age.

Please see additional Important Safety Information throughout and the accompanying full Prescribing Information.



Not actual size.

About psoriasis

What is plaque psoriasis?

- Plaque psoriasis is the most common form of psoriasis, but the exact cause is unknown
- This condition leads to your immune system influencing your skin cells to develop too quickly
- Rapid development of cells causes immature skin cells to build up into red, scaly, thickened patches of skin called plaques
- Symptoms can include burning, skin pain, cracking of the skin, and itching

What triggers plaque psoriasis?

What causes flares?

There are certain triggers that cause plaque psoriasis to flare. Some of the most common triggers include:



Bacterial infections



Medications



Skin injuries



Smoking



Stress



Not an actual patient.

Symptoms of plaque psoriasis may vary.

TALK WITH YOUR DOCTOR TO LEARN MORE.

Proven effective treatment for plaque psoriasis

How to use Enstilar® Spray Foam

Wash your hands before applying Enstilar® Spray Foam.



Not actual size.

- ✓ A majority of patients (53%) were “Clear” or “Almost Clear” at Week 4 in a clinical trial
- ✓ **Rapid results** may occur at Week 2
- ✓ **Once-daily application** for up to 4 weeks



SHAKE

the Enstilar® Spray Foam can before each application.



SPRAY

the foam at least 1.5 (one-and-a-half) inches away from the skin by pushing down the nozzle, in any position, except horizontally (sideways).



RUB

Enstilar® Spray Foam gently on an affected body area until medication is absorbed. After applying Enstilar® Spray Foam, put the cap back on and wash your hands.

IMPORTANT SAFETY INFORMATION *(cont'd)*

Enstilar Foam is for use on skin only (topical). Do not get Enstilar Foam near or in your mouth, eyes, or vagina. Avoid using Enstilar Foam on your face, groin, or armpits, or if you have thinning of your skin (atrophy) at the treatment site.

Please see additional Important Safety Information throughout and the [accompanying](#) full Prescribing Information.

Repeat these 3 steps to apply Enstilar® Spray Foam to other affected areas as instructed by your doctor.

Results achieved in clinical trials

The photos on this page are actual patient images from Enstilar® Spray Foam clinical trials. These images represent before treatment, at Week 1, and at Week 4.

Individual results may vary.



Not actual size.

ARM

LEG

ELBOW

KNEE

BEFORE treatment



WEEK 1 of treatment



WEEK 4 of treatment



IMPORTANT SAFETY INFORMATION (cont'd)

Enstilar Foam is flammable. Avoid fire, flame, and smoking when applying and right after you apply Enstilar Foam.

Enstilar Foam has the appearance of a non-expanding foam that gradually collapses after spraying.

Please see additional Important Safety Information throughout and the accompanying full Prescribing Information.

Enstilar®
(calcipotriene and betamethasone dipropionate) Foam 0.005%/0.064%

Save on your Enstilar Spray Foam® prescription with the LEO Pharma® CONNECT Savings Program

The LEO Pharma® CONNECT program is a company-sponsored patient access and support program that is available to assist commercially insured patients.

2 steps

TO SAVINGS FOR
ELIGIBLE COMMERCIALY
INSURED PATIENTS

STEP 1

Activate an existing card or request a card by calling **1-888-727-6325** or by visiting **leopharmaconnect.com**

STEP 2

Present the card with your prescription at any pharmacy

*Certain restrictions apply. The LEO Pharma® CONNECT program may reduce out-of-pocket expenses. Must be 12 years of age or older to be eligible, and a legal guardian over 18 years of age must redeem the card for patients aged 12 to 17. You are not eligible if you are enrolled or you participate in any state or federally funded health care program (eg, Medicare, Medicaid, etc). Full details of the LEO Pharma® CONNECT program are available at leopharmaconnect.com or may be obtained by calling 1-877-678-7494 between 8:30 AM and 8:30 PM (Eastern), Monday through Friday.

Enstilar®
(calcipotriene and betamethasone dipropionate) Foam 0.005%/0.064%

LEO Pharma®
CONNECT
Access. Affordability. Assistance.

Most eligible patients **\$20***
pay as little as **\$20***
per prescription of Enstilar® Spray Foam



Not an actual
co-pay card.

IMPORTANT SAFETY INFORMATION *(cont'd)*

What should I discuss with my healthcare provider before starting Enstilar Foam?

Tell your healthcare provider about all your medical conditions, including if you:

- have a calcium metabolism disorder
- have thinning skin (atrophy) at the treatment site
- are pregnant or plan to become pregnant. It is not known if Enstilar Foam will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Enstilar Foam passes into your breastmilk.

Tell your healthcare provider about all the medicines you take, including any corticosteroid medicines, any other products containing calcipotriene, and any other prescription and over-the-counter medicines, vitamins and herbal supplements. Do not use other products containing calcipotriene or a corticosteroid medicine without talking to your healthcare provider first.

How should I use Enstilar Foam?

Read the Instructions For Use that comes with Enstilar Foam. There you will find information about how to apply and store Enstilar Foam.

Use Enstilar Foam exactly as your healthcare provider says.

Do not use more than 60 grams of Enstilar Foam every 4 days.

Do not use Enstilar Foam for longer than prescribed. Using too much Enstilar Foam or using it too often or for too long can increase your risk for having serious side effects.

Do not bandage or cover the treated skin area, unless instructed by your healthcare provider.

Gently rub in Enstilar Foam to your affected areas. Wash your hands after applying Enstilar Foam unless you are using it to treat your hands.

Apply Enstilar Foam to the affected areas of your skin 1 time a day for up to 4 weeks. You should stop treatment when your plaque psoriasis is under control, unless your healthcare provider gives you other instructions.

Enstilar[®]

(calcipotriene and betamethasone dipropionate) Foam 0.005%/0.064%

What are the possible side effects of Enstilar Foam?

Enstilar Foam may cause serious side effects including:

- **Too much calcium in your blood or urine and/or adrenal gland problems**
- **Cushing's syndrome**, a condition that happens when your body is exposed to too much of the hormone cortisol
- **High blood sugar and sugar in your urine**
- **Skin problems, including reactions where Enstilar Foam is applied, and allergic reactions** (allergic contact dermatitis). Tell your healthcare provider if you have any skin problems including thinning of your skin, pain/burning, inflammation, itching, irritation, dryness, changes in skin color, redness, infection or raised bumps on your skin.
- **Eye problems:** Enstilar Foam may increase your chance of getting cataracts and glaucoma. Do not get Enstilar Foam in your eyes because it may cause irritation. If you accidentally get Enstilar Foam on your face, in your mouth or in your eyes, wash the area with water right away. Tell your healthcare provider if you have blurred vision or get any other vision problems during treatment.

The most common side effects include irritation, itching, inflamed hair pores (folliculitis), changes in skin color, rash with raised red bumps or skin welts (hives) and worsening of your psoriasis. These are not all the possible side effects. Call your healthcare provider for medical advice about side effects.

Where do I learn more?

This summary does not include all the information about Enstilar Foam. Read the information that comes with your prescription every time you have a prescription filled. Talk to your healthcare provider or pharmacist about this information. Your healthcare provider is the best person to help you decide if Enstilar Foam is right for you.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

You may also report side effects to LEO Pharma Inc. at 1-877-494-4536, option 1, or email to usdrugsafety@leo-pharma.com.

Please see additional Important Safety Information throughout and the [accompanying full Prescribing Information](#).

Enstilar[®]

(calcipotriene and betamethasone dipropionate) Foam 0.005%/0.064%

Clearer skin is possible in just 4 weeks

✓ Proven effective to treat plaque psoriasis

A majority of patients (53%) were “Clear” or “Almost Clear” at Week 4 in a clinical trial

✓ Spray-on foam

A once-daily spray foam that can be applied **across a large body area**

(do not use more than 60 grams of Enstilar[®] Spray Foam every 4 days)



Not actual size.

**BE SURE TO DISCUSS
YOUR TREATMENT GOALS
WITH YOUR DOCTOR**

IMPORTANT SAFETY INFORMATION *(cont'd)*

Enstilar Foam is for use on skin only (topical). Do not get Enstilar Foam near or in your mouth, eyes, or vagina. Avoid using Enstilar Foam on your face, groin, or armpits, or if you have thinning of your skin (atrophy) at the treatment site.

Please see additional Important Safety Information throughout and the accompanying full Prescribing Information.



The LEO Pharma logo, LEO Pharma, and Enstilar are registered trademarks of LEO Pharma A/S. ©2021 LEO Pharma Inc. All rights reserved. March 2021 MAT-42142